Geometric Modeling

Summer Semester 2012

Differential Geometry

(and a bit of Topology)







Multi-Dimensional Derivatives

Derivative of a Function

Reminder: The *derivative* of a function is defined as

- $\frac{d}{dt}f(t) \coloneqq \lim_{h \to 0} \frac{f(t+h) f(t)}{h}$
- **If limit exists:** function is called *differentiable*.

Other notation:

 $\frac{d}{dt}f(t) = \underbrace{f'(t)}_{\text{variable}} = \underbrace{\dot{f}(t)}_{\text{time}}$

from context

variables

 $\frac{d^k}{dt^k}f(t)=f^{(k)}(t)$

repeated differentiation (higher order derivatives)

Taylor Approximation

Smooth functions can be approximated locally:

$$f(x) \approx f(x_0) + \frac{d}{dx} f(x_0) (x - x_0) + \frac{1}{2} \frac{d^2}{dx^2} f(x_0) (x - x_0)^2 + \dots \dots + \frac{1}{k!} \frac{d^k}{dx^k} f(x_0) (x - x_0)^k + O(x^{k+1})$$

- Convergence: holomorphic functions
- Local approximation for smooth functions

Rule of Thumb

Derivatives and Polynomials

- Polynomial: $f(x) = c_0 + c_1 x + c_2 x^2 + c_3 x^3 \dots$
 - Oth-order derivative: $f(0) = c_0$
 - 1st-order derivative: $f'(0) = c_1$
 - 2nd-order derivative: $f''(0) = 2c_2$
 - 3rd-order derivative: $f'''(0) = 6c_3$

• ...

Rule of Thumb:

- Derivatives correspond to polynomial coefficients
- Estimate derivatives ↔ polynomial fitting

Partial Derivative

Multivariate functions:

• Notation changes:

$$\frac{\partial}{\partial x_{k}} f(x_{1}, ..., x_{k-1}, x_{k}, x_{k+1}, ..., x_{n}) := \lim_{h \to 0} \frac{f(x_{1}, ..., x_{k-1}, x_{k} + h, x_{k+1}, ..., x_{n}) - f(x_{1}, ..., x_{k-1}, x_{k}, x_{k+1}, ..., x_{n})}{h}$$

• Alternative notation:

$$\frac{\partial}{\partial x_k} f(\mathbf{x}) = \partial_k f(\mathbf{x}) = f_{x_k}(\mathbf{x})$$

Special Cases

Derivatives for:

- Functions $f: \mathbb{R}^n \to \mathbb{R}$ ("heightfield")
- Functions $f: \mathbb{R} \to \mathbb{R}^n$ ("curves")
- Functions $f: \mathbb{R}^n \to \mathbb{R}^m$ (general case)

Special Cases

Derivatives for:

- Functions $f: \mathbb{R}^n \to \mathbb{R}$ ("height field")
- Functions $f: \mathbb{R} \to \mathbb{R}^n$ ("curves")
- Functions $f: \mathbb{R}^n \to \mathbb{R}^m$ (general case)



height field

Gradient

Gradient:

- Given a function $f: \mathbb{R}^n \to \mathbb{R}$ ("heightfield")
- The vector of all partial derivatives of *f* is called the *gradient*:

$$\nabla f(\mathbf{x}) = \begin{pmatrix} \frac{\partial}{\partial x_1} \\ \vdots \\ \frac{\partial}{\partial x_n} \end{pmatrix} f(\mathbf{x}) = \begin{pmatrix} \frac{\partial}{\partial x_1} f(\mathbf{x}) \\ \vdots \\ \frac{\partial}{\partial x_n} f(\mathbf{x}) \end{pmatrix}$$

Gradient

Gradient:



- gradient: vector pointing in direction of steepest ascent.
- Local linear approximation (Taylor):

 $f(\mathbf{x}) \approx f(\mathbf{x}_0) + \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0)$

Higher Order Derivatives

Higher order Derivatives:

• Can do all combinations:

$$\left(\frac{\partial}{\partial x_{i_1}}\frac{\partial}{\partial x_{i_2}}\dots\frac{\partial}{\partial x_{i_k}}\right)f$$

• Order does not matter for $f \in C^k$

Hessian Matrix

Higher order Derivatives:

• Important special case: Second order derivative

$$\begin{pmatrix} \frac{\partial^2}{\partial x_1^2} & \frac{\partial}{\partial x_2} \frac{\partial}{\partial x_1} & \cdots & \frac{\partial}{\partial x_n} \frac{\partial}{\partial x_1} \\ \frac{\partial}{\partial x_1} \frac{\partial}{\partial x_2} & \frac{\partial^2}{\partial x_1^2} & \cdots & \frac{\partial}{\partial x_n} \frac{\partial}{\partial x_2} \\ \vdots & \ddots & \vdots \\ \frac{\partial}{\partial x_1} \frac{\partial}{\partial x_n} & \frac{\partial}{\partial x_2} \frac{\partial}{\partial x_n} & \cdots & \frac{\partial^2}{\partial x_n^2} \end{pmatrix} f(\mathbf{x}) =: H_f(\mathbf{x})$$

- "Hessian" matrix (symmetric for $f \in C^2$)
- Orthogonal Eigenbasis, full Eigenspectrum

Taylor Approximation



2nd order approximation (schematic)

Second order Taylor approximation:

• Fit a paraboloid to a general function

 $f(\mathbf{x}) \approx f(\mathbf{x}_0) + \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0) + \frac{1}{2} (\mathbf{x} - \mathbf{x}_0)^{\mathrm{T}} \cdot H_f(\mathbf{x}_0) \cdot (\mathbf{x} - \mathbf{x}_0)$

Special Cases

Derivatives for:

- Functions $f: \mathbb{R}^n \to \mathbb{R}$ ("heightfield")
- Functions $f: \mathbb{R} \to \mathbb{R}^n$ ("curves")
- Functions $f: \mathbb{R}^n \to \mathbb{R}^m$ (general case)



curves

Derivatives of Curves

Derivatives of vector valued functions:

• Given a function $f: \mathbb{R} \to \mathbb{R}^n$ ("curve")

 $f(t) = \begin{pmatrix} f_1(t) \\ \vdots \\ f_n(t) \end{pmatrix}$

• We can compute derivatives for every output dimension:

$$\frac{d}{dt}f(t) \coloneqq \begin{pmatrix} \frac{d}{dt}f_1(t) \\ \vdots \\ \frac{d}{dt}f_n(t) \end{pmatrix} \coloneqq f'(t) \rightleftharpoons \dot{f}(t)$$

Geometric Meaning



Tangent Vector:

- f': tangent vector
- Motion of physical particle: \dot{f} = velocity.
- Higher order derivatives: Again vector functions
- Second derivative \ddot{f} = acceleration

Special Cases

Derivatives for:

- Functions $f: \mathbb{R}^n \to \mathbb{R}$ ("heightfield")
- Functions $f: \mathbb{R} \to \mathbb{R}^n$ ("curves")
- Functions $f: \mathbb{R}^n \to \mathbb{R}^m$ (general case)



You can combine it...

General case:

• Given a function $f: \mathbb{R}^n \to \mathbb{R}^m$ ("space warp")

 $f(\mathbf{x}) = f((x_1, ..., x_n)) = \begin{pmatrix} f_1(x_1, ..., x_n) \\ \vdots \\ f_m(x_1, ..., x_n) \end{pmatrix}$

- Maps points in space to other points in space
- First derivative: Derivatives of all *output components* of *f* w.r.t. all *input directions*.
- "Jacobian matrix": denoted by ∇f or \mathbf{J}_f

Jacobian Matrix

Jacobian Matrix:

$$\nabla f(\mathbf{x}) = J_f(\mathbf{x}) = \nabla f(x_1, \dots, x_n)^{\mathrm{T}}$$
$$= \begin{pmatrix} \nabla f_1(x_1, \dots, x_n)^{\mathrm{T}} \\ \vdots \\ \nabla f_m(x_1, \dots, x_n)^{\mathrm{T}} \end{pmatrix} = \begin{pmatrix} \partial_{x_1} f_1(\mathbf{x}) & \cdots & \partial_{x_n} f_1(\mathbf{x}) \\ \vdots \\ \partial_{x_1} f_m(\mathbf{x}) & \cdots & \partial_{x_n} f_m(\mathbf{x}) \end{pmatrix}$$

Use in a first-order Taylor approximation:

$$f(\mathbf{x}) \approx f(\mathbf{x}_0) + J_f(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0)$$

$$\uparrow$$
matrix / vector

product

Special Cases

Derivatives for:

- Functions $f: \mathbb{R}^n \to \mathbb{R}$ ("heightfield")
- Functions $f: \mathbb{R} \to \mathbb{R}^n$ ("curves")
- Functions $f: \mathbb{R}^n \to \mathbb{R}^m$ (general case)



Coordinate Systems

Problem:

- What happens, if the coordinate system changes?
- Partial derivatives go into different directions then.
- Do we get the same result?

Total Derivative

First order Taylor approx.:

- $f(\mathbf{x}_0) + \nabla f(\mathbf{x}_0) \cdot (\mathbf{x} \mathbf{x}_0) + R_{x_0}(\mathbf{x})$
- Converges for C¹ functions $f: \mathbb{R}^n \to \mathbb{R}^m$

$$\lim_{\mathbf{x}\to\mathbf{x}_0}\frac{R_{x_0}(\mathbf{x})}{\|\mathbf{x}-\mathbf{x}_0\|}=0,$$

("totally differentiable")



Consequences:

- A linear function: fully determined by image of a basis
- Hence: Directions of partial derivatives do not matter this is just a basis transform.
 - We can use any linear independent set of directions T
 - Transform to standard basis by multiplying with T⁻¹
- Similar argument for higher order derivatives

Directional Derivative

The directional derivative is defined as:

- Given $f: \mathbb{R}^n \to \mathbb{R}^m$ and $\mathbf{v} \in \mathbb{R}^n$, $||\mathbf{v}|| = 1$.
- Directional derivative:

$$\nabla_{\mathbf{v}} f(\mathbf{x}) = \frac{\partial f}{\partial \mathbf{v}}(\mathbf{x}) \coloneqq \frac{d}{dt} f(\mathbf{x} + t\mathbf{v})$$

• Compute from Jacobian matrix

$$\nabla_{\mathbf{v}} f(\mathbf{x}) = \nabla f(\mathbf{x}) \frac{\mathbf{v}}{\|\mathbf{v}\|}$$

(requires total differentiability)

Multi-Dimensional Integrals

Integral

Integral of a function

- Function $f: \mathbb{R} \to \mathbb{R}$
- Integral $\int_{a}^{b} f(t)dt$ measures signed area under curve:



Integral

Numerical Approximation

• Sum up a series of approximate shapes



• (Riemannian) Definition: limit for baseline \rightarrow zero

Multi-Dimensional Integral

Integration in higher dimensions

- Functions $f: \mathbb{R}^n \to \mathbb{R}$
- Tessellate domain and sum up volume of cuboids



Integral Transformations

Integration by substitution:

 $\int_{a}^{b} f(x) dx = \int_{g^{-1}(a)}^{g^{-1}(b)} f(g(t))g'(t) dt$

Need to compensate for speed of movement that shrinks the measured area.



Multi-Dimensional Substitution

Transformation of Integrals:

$$\int_{\Omega} f(\mathbf{x}) d\mathbf{x} = \int_{g^{-1}(\Omega)} f(g(\mathbf{y})) |\det(\nabla g(\mathbf{y}))| d\mathbf{y}$$

- $g \in C^1$, invertible
- Jacobian approximates local behavior of g()



- Determinant: local area/volume change
- In particular: $|\det(\nabla g(\mathbf{y}))| = 1$ means g() is *area/volume conserving*.

Topology - a very short primer -

A Few Concepts from Topology

Homeomorphism:

- $f: X \to Y$
- *f* is bijective
- *f* is continuous
- f⁻¹ exists and is continuous
- Basically, a continuous deformation

Topological equivalence

- Objects are topologically equivalence if there exists a homeomorphism that maps between them
- "Can be deformed into each other"



Surfaces

Boundaries of volumes in 3D

- Topological Equivalence classes
 - Sphere
 - Torus
 - n-fold Torus
- Genus = number of tunnels



Manifold

Definition: Manifold

• A *d*-manifold *M*:

At every $x \in M$ there exists an ϵ -environment homeomorphic to a *d*-dimensional disc

• With boundary: *disc* or *half-disc*

Further concepts

Connected Set

 There exists a continuous curve within the set between all pairs of points

Simply Connected

 Every closed loop can be continuously shrunken until it disappears

Differential Geometry of Curves & Surfaces
Part I: Curves

Parametric Curves

Parametric Curves:

• A differentiable function

 $f: (a, b) \rightarrow \mathbb{R}^n$



describes a *parametric curve* $C = f((a, b)), C \subseteq \mathbb{R}^{n}.$

- The parametrization is called *regular* if $f'(t) \neq 0$ for all t.
- If || f'(t) || = 1 for all t, f is called a unit-speed parametrization of the curve C.

Length of a Curve

The length of a curve:

- The length of a regular curve *C* is defined as: length_C = $\int_{a}^{b} ||f'(t)|| dt$
- Independent of the parametrization (integral transformation theorem).
- Alternative: $length_c = |b a|$ for a unit-speed parametrization

Reparametrization

Enforcing unit-speed parametrization:

- Assume: $||f'(t)|| \neq 0$ for all t.
- We have:

length_c(x) = $\int_{a}^{x} ||f'(t)|| dt$ (invertible, because f'(t) > 0)

Concatenating f
 length⁻¹ yields a unit-speed

 parametrization of the curve



Tangents

Unit Tangents:

• The unit tangent vector at $x \in (a, b)$ is given by:

 $\operatorname{tangent}(t) = \frac{f'(t)}{\|f'(t)\|}$

For curves C ⊂ ℝ², the unit normal vector of the curve is defined as:

normal(t) =
$$\begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \frac{f'(t)}{\|f'(t)\|}$$

Curvature

Curvature:

- First derivatives: curve direction / speed.
- Curvature encoded in 2nd order information.
- Why not just use *f*"?
- Problem: Depends on parametrization
 - Different velocity yields different results
 - Need to distinguish between acceleration in tangential and non-tangential directions.

Curvature & 2nd Derivatives



Definition of curvature

- We want only the non-tangential component of f''.
- Accelerating/slowing down does not matter for curvature of the traced out curve C.
- Need to normalize speed.

Curvature

Curvature of a Curve $C \in \mathbb{R}^2$:

$$\kappa \mathbf{2}(t) = \frac{\left\langle f''(t), \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} f'(t) \right\rangle}{\left\| f'(t) \right\|^3}$$

- Normalization factor:
 - Divide by ||f'|| to obtain unit tangent vector
 - Divide again twice to normalize f"
 - Taylor expansion / chain rule:

$$f(\lambda t) = f(t_0) + \lambda f'(t_0)(t - t_0) + \frac{1}{2}\lambda^2 f''(t)(t - t_0)^2 + O(t^3)$$

Second derivative scales quadratically with speed

Unit-speed parametrization

Unit-speed parametrization:

- Assume a unit-speed parametrization, i.e. ||f'|| = 1.
- Then, **κ**2 simplifies to:

 $\mathbf{\kappa}\mathbf{2}(t) = \left\| f^{\prime\prime}(t) \right\|$

Radius of Curvature

Easy to see:

- Curvature of a circle is constant, $\kappa 2 \equiv \pm 1/r$ (r = radius).
- Accordingly: Define radius of curvature as $1/\kappa 2$.
- Osculating circle:
 - Radius: 1/κ2
 - Center: $f(t) + \frac{1}{\kappa^2} normal(t)$



Theorems

Definition:

- Rigid motion: $\mathbf{x} \rightarrow \mathbf{Ax+b}$ with orthogonal **A**
 - Orientation preserving (no mirroring) if det(A) = +1
 - Mirroring leads to det(A) = -1

Theorems for plane curves:

- Curvature is invariant under rigid motion
 - Absolute value is invariant
 - Signed value is invariant for orientation preserving rigid motion
- Two unit speed parameterized curves with identical signed curvature function differ only in a orientation preserving rigid motion.

Space Curves

General case: Curvature of a Curve $C \subseteq \mathbb{R}^n$

- Given a unit-speed parametrization *f* of C
- The *curvature* of C at parameter value *t* is defined as: $\kappa(t) = ||f''(t)||$
- For a general, regular curve C ⊆ ℝ³ (any regular parametrization):

 $\mathbf{\kappa}(t) = \frac{\left\|f'(t) \times f''(t)\right\|}{\left\|f'(t)\right\|^3}$

General curvature is unsigned



Torsion

Characteristics of Space Curves in \mathbb{R}^3 :

- Curvature not sufficient
- Curve may "bend" in space
- Curvature is a 2nd order property
- 2nd order curves are always flat
 - Quadratic curves are specified by 3 points in space, which always lie in a plane
 - Cannot capture out-of-plane bends
- Missing property: Torsion

Torsion

Definition:

- Let f be a regular parametrization of a curve $C \subseteq \mathbb{R}^3$ with non-zero curvature
- The torsion of *f* at *t* is defined as

 $\boldsymbol{\tau}(t) = \frac{f'(t) \times f''(t) \cdot f'''(t)}{\|f'(t) \times f''(t)\|^2} = \frac{\det(f'(t), f''(t), f'''(t))}{\|f'(t) \times f''(t)\|^2}$

Illustration



Theorem

Fundamental Theorem of Space Curves

• Two unit speed parameterized curves $C \subseteq \mathbb{R}^3$ with identical, positive curvature and identical torsion are identical up to a rigid motion.

Part II: Surfaces

Parametric Patches

Parametric Surface Patches:

A smoothly differentiable function

 $f: \mathbb{R}^2 \supseteq \Omega \to \mathbb{R}^n$

describes a *parametric surface patch*

 $\mathsf{P} = f(\Omega), \, \mathsf{P} \subseteq \mathbb{R}^n.$

Parametric Patches

Function $f(\mathbf{x}) = f(u, v) \to \mathbb{R}^3$

- Tangents: $\frac{d}{dt}f(\mathbf{x}_0 + t\mathbf{r}) = \nabla_{\mathbf{r}}f(\mathbf{x}_0)$
- Canonical tangents: $\partial_u f(u,v), \partial_v f(u,v)$
- Normal: $\mathbf{n}(\mathbf{x}_0) = \frac{\partial_u f(u,v) \times \partial_v f(u,v)}{\|\partial_u f(u,v) \times \partial_v f(u,v)\|}$

Illustration



Surface Area

Surface Area:

- Patch $P: f: \Omega \to \mathbb{R}^3$
- Computation is simple
- Integrate over constant function $f \equiv 1$ over surface
- Then apply integral transformation theorem:

area(P) =
$$\int_{\Omega} \|\partial_u f(\mathbf{x}) \times \partial_v f(\mathbf{x})\| d\mathbf{x}$$

Fundamental Forms

Fundamental Forms:

- Describe the local parametrized surface
- Measure...
 - ...distortion of length (first fundamental form)
 - ...surface curvature (second fundamental form)
- Parametrization independent surface curvature measures will be derived from this

First Fundamental Form

First Fundamental Form

- Also known as *metric tensor*.
- Given a regular parametric patch $f: \mathbb{R}^2 \supseteq \Omega \rightarrow \mathbb{R}^3$.
- *f* will distort angles and distances
- We will look at a local first order Taylor approximation to measure the effect:

 $f(\mathbf{x}) \approx f(\mathbf{x}_0) + \nabla f(\mathbf{x}_0) (\mathbf{x} - \mathbf{x}_0)$

• Length changes become visible in the scalar product...



First Fundamental Form

First Fundamental Form

- First order Taylor approximation: $f(\mathbf{x}) \approx f(\mathbf{x}_0) + \nabla f(\mathbf{x}_0)(\mathbf{x} - \mathbf{x}_0)$
- Scalar product of vectors $\mathbf{a}, \mathbf{b} \in \mathbb{R}^2$: $\langle f(\mathbf{x}_0 + \mathbf{a}) - f(\mathbf{x}_0), f(\mathbf{x}_0 + \mathbf{b}) - f(\mathbf{x}_0) \rangle \approx \langle \nabla f(\mathbf{x}_0) \mathbf{a}, \nabla f(\mathbf{x}_0) \mathbf{b} \rangle$ $= \mathbf{a}^{\mathrm{T}} (\nabla f(\mathbf{x}_0)^{\mathrm{T}} \nabla f(\mathbf{x}_0)) \mathbf{b}$

first fundamental form

 $\partial_{v} f(\mathbf{x}_{0})$

 \mathbf{X}_0 U $f(\mathbf{x}_0)$

First Fundamental Form

First Fundamental Form

The first fundamental form can be written as a 2 × 2 matrix:

 $\left(\nabla f^{\mathrm{T}} \nabla f \right) = \begin{pmatrix} \partial_{\mathbf{u}} f \partial_{\mathbf{u}} f & \partial_{\mathbf{u}} f \partial_{\mathbf{v}} f \\ \partial_{\mathbf{u}} f \partial_{\mathbf{v}} f & \partial_{\mathbf{v}} f \partial_{\mathbf{v}} f \end{pmatrix} = : \begin{pmatrix} E & F \\ F & G \end{pmatrix} \qquad \mathbf{I}(\mathbf{x}, \mathbf{y}) := \mathbf{x}^{\mathrm{T}} \left(\nabla f^{\mathrm{T}} \nabla f \right) \mathbf{y}$

- The matrix is symmetric and positive definite (regular parametrization, semi-definite otherwise)
- Defines a *generalized scalar product* that measures lengths and angles *on the surface*.

Second Fundamental Form

Problems:

- The first fundamental form measures length changes only.
- A cylinder looks like a flat sheet in this view.
- We need a tool to measure curvature of a surface as well.
- This requires second order information.
 - Any first order approximation is inherently "flat".

Second Fundamental Form

Definition:

- Given: regular parametric patch $f: \mathbb{R}^2 \supseteq \Omega \rightarrow \mathbb{R}^3$.
- Second fundamental form:

(a.k.a. shape operator, curvature tensor)

$$S(\mathbf{x}_0) = \begin{pmatrix} \partial_{\mathbf{u}\mathbf{u}} f(\mathbf{x}_0) \cdot \mathbf{n} & \partial_{\mathbf{u}\mathbf{v}} f(\mathbf{x}_0) \cdot \mathbf{n} \\ \partial_{\mathbf{u}\mathbf{v}} f(\mathbf{x}_0) \cdot \mathbf{n} & \partial_{\mathbf{v}\mathbf{v}} f(\mathbf{x}_0) \cdot \mathbf{n} \end{pmatrix}$$

• Notation:

$$\mathbf{II}(\mathbf{x},\mathbf{y}) = \mathbf{x}^{\mathrm{T}} \begin{pmatrix} \partial_{\mathbf{u}\mathbf{u}} f(\mathbf{x}_{0}) \cdot \mathbf{n} & \partial_{\mathbf{u}\mathbf{v}} f(\mathbf{x}_{0}) \cdot \mathbf{n} \\ \partial_{\mathbf{u}\mathbf{v}} f(\mathbf{x}_{0}) \cdot \mathbf{n} & \partial_{\mathbf{v}\mathbf{v}} f(\mathbf{x}_{0}) \cdot \mathbf{n} \end{pmatrix} \mathbf{y}$$

Second Fundamental Form

Basic Idea:

- Compute second derivative vectors
- Project in normal direction (remove tangential acceleration)

Alternative Computation

Alternative Formulation (Gauss):

- Local height field parameterization f(x,y) = z
- Orthonormal x,y coordinates tangential to surface,
 z in normal direction, origin at zero

Ζ

X

• 2nd order Taylor representation:

$$f(\mathbf{x}) \approx \frac{1}{2} \underbrace{\mathbf{x}^{\mathrm{T}} f''(\mathbf{x}) \mathbf{x}}_{2} + \underbrace{f'(\mathbf{x}) \mathbf{x} + f(0)}_{0}$$
$$= ex^{2} + 2fxy + gy^{2}$$

• Second fundamental form: Matrix of second derivatives

$$\begin{pmatrix} \partial_{xx} f & \partial_{xy} f \\ \partial_{xy} f & \partial_{yy} f \end{pmatrix} = \begin{pmatrix} e & f \\ f & g \end{pmatrix}$$

Basic Idea

In other words:

- First fundamental form: I Linear part (squared) of local Taylor approximation.
- *Second fundamental form:* **II** Quadratic part of heightfield approximation
- Both matrices are symmetric.
 - Next: eigenanalysis, of course...



Principal Curvature

Eigenanalysis:

- Eigenvalues of second fundamental form for an orthonormal tangent basis are called principal curvatures κ₁, κ₂.
- Corresponding orthogonal eigenvectors are called principal directions of curvature.



Normal Curvature

Definition:

• The *normal curvature* $k(\mathbf{r})$ in direction \mathbf{r} for a unit length direction vector \mathbf{r} at parameter position \mathbf{x}_0 is given by: $k_{\mathbf{x}_0}(\mathbf{r}) = \mathbf{II}_{\mathbf{x}_0}(\mathbf{r},\mathbf{r}) = \mathbf{r}^{\mathrm{T}}\mathbf{S}(\mathbf{x}_0)\mathbf{r}$

Relation to Curvature of Plane Curves:

- Intersect the surface locally with plane spanned by normal and r through point x₀.
- Identical curvatures (up to sign).



Principal Curvatures

Relation to principal curvature:

- The maximum principal cuvature κ_1 is the maximum of the normal curvature
- The minimum principal cuvature $\kappa_{\rm 2}$ is the minimum of the normal curvature

Gaussian & Mean Curvature

More Definitions:

- The Gaussian curvature K is the product of the principal curvatures: $K = \kappa_1 \kappa_2$
- The mean curvature *H* is the average: $H = 0.5 \cdot (\kappa_1 + \kappa_2)$

Theorems:

•
$$K(\mathbf{x}_0) = \det(S(x_0)) = \frac{eg - f^2}{EG - F^2}$$

• $H(\mathbf{x}_0) = \frac{1}{2} \operatorname{tr}(S(x_0)) = \frac{eG - 2fF + gE}{2(EG - F^2)}$

Global Properties

Definition:

- Isometry: mapping between surfaces that preserves distances on the surface (geodesics)
- Developable surface: Gaussian curvature zero everywhere (i.e. no curvature in at least one direction)
 - Examples: Cylinder, Cone, Plane
- A developable surface can be locally mapped to a plane isometrically (flattening out, unroll).

Theorema Egregium

Theorema egregium (Gauss):

- Any isometric mapping preservers Gaussian curvature, i.e. Gaussian curvature is invariant under isometric maps ("intrinsic surface property")
- Consequence: The earth (≈ sphere) cannot be mapped to a plane in an exactly length preserving way.
Gauss Bonnet Theorem

Gauss Bonnet Theorem:

For a compact, orientable surface without boundary in \mathbb{R}^3 , the area integral of the Gauss curvature is related to the genus g of the surface:

$$\int_{S} K(x) dx = 4 \pi (1 - g)$$



Fundamental Theorem of Surfaces

Theorem:

- Given two parametric patches in \mathbb{R}^3 defined on the same domain Ω .
- Assume that the first and second fundamental form are identical.
- Then there exists a rigid motion that maps on surface to the other.

Summary

Objects are the same up to a rigid motion, if...:

- Curves $\mathbb{R} \to \mathbb{R}^2$: Same *speed*, same *curvature*
- Curves $\mathbb{R} \to \mathbb{R}^3$: Same *speed*, same *curvature*, *torsion*
- Surfaces $\mathbb{R}^2 \to \mathbb{R}^3$: Same *first* & *second* fundamental form
- Volumetric Objects $\mathbb{R}^3 \rightarrow \mathbb{R}^3$: Same *first* fundamental form



Deformation Models

What if this does not hold?

- Deviation in fundamental forms is a measure of deformation
- Example: Surfaces
 - Diagonals of I₁ I₂: scaling (stretching)
 - Off-diagonals of I₁ I₂: sheering
 - Elements of II₁ II₂: bending
- This is the basis of *deformation models*.

Reference: D. Terzopoulos, J. Platt, A. Barr, K. Fleischer: Elastically Deformable Models. In: *Siggraph '87 Conference Proceedings (Computer Graphics 21(4))*, 1987.



Part III: Intrinsic Geometry and Intrinsic Mappings

Scenario

Mapping between Surfaces

- Intrinsic view only metric tensor
- Ignore isometric deformations
- Applications:
 - Deformable shape matching
 - Texture mapping (flat → 3D) "parametrization"



 S_2

Riemannian Manifolds

Concept:

- Abstract surface
- "User defined" metric



Riemannian Manifolds

Riemannian Manifold

- Manifold topology, *d*-dimensional
 - We mostly focus on 2-manifolds, embedded in \mathbb{R}^3
- Local parametrization: tangent space
 - We have a tangent space for each point
- Intrinsic metric
 - For every point, we know the metric tensor

 (i.e., a scalar product)
 .
 - In tangent space coordinates



Types of Mappings

Given: Riemannian Manifolds M₁, M₂

Consider: Functions $f: M_1 \rightarrow M_2$

Important types of mappings:

- Isometric: preserves distances & angles
 - Jacobian: $\nabla f = \mathbf{R}$, \mathbf{R} orthogonal
 - Metric tensor: I
- Conformal: preserves (only) angles
 - Jacobian: $\nabla f = \lambda \mathbf{R}, \lambda > 0, \mathbf{R}$ orthogonal
 - Metric tensor: $\lambda \mathbf{I}, \lambda > 0$
- Equi-areal (incompressible): preserves area
 - Jacobian: $|\det(\nabla f)| = 1$
 - Metric tensor: det(M) = 1







Theorems

Isometric mappings

Isometries preserve Gaussian curvature (Gauss)

Conformal mappings

- Uniformization theorem:
 - For any simply connected Riemann surface (2-manifold)
 - There exists a conformal, bijective mapping to either:
 - the open unit disk
 - the whole 2D plane
 - Unit sphere
- Any discs, spheres, planes can be mapped to objects of the same topology using a conformal map