

3.10 Refutational Completeness of Resolution

How to show refutational completeness of propositional resolution:

- We have to show: $N \models \perp \Rightarrow N \vdash_{Res} \perp$, or equivalently: If $N \not\vdash_{Res} \perp$, then N has a model.
- Idea: Suppose that we have computed sufficiently many inferences (and not derived \perp).
- Now order the clauses in N according to some appropriate ordering, inspect the clauses in ascending order, and construct a series of Herbrand interpretations.
- The limit interpretation can be shown to be a model of N .

Clause Orderings

1. We assume that \succ is any fixed ordering on ground atoms that is *total* and *well-founded*. (There exist many such orderings, e.g., the length-based ordering on atoms when these are viewed as words over a suitable alphabet.)
2. Extend \succ to an *ordering* \succ_L on *ground literals*:

$$\begin{array}{l} [\neg]A \succ_L [\neg]B \quad , \text{ if } A \succ B \\ \neg A \succ_L A \end{array}$$

3. Extend \succ_L to an *ordering* \succ_C on *ground clauses*:
 $\succ_C = (\succ_L)_{mul}$, the multi-set extension of \succ_L .
Notation: \succ also for \succ_L and \succ_C .

Example

Suppose $A_5 \succ A_4 \succ A_3 \succ A_2 \succ A_1 \succ A_0$. Then:

$$\begin{array}{l} A_0 \vee A_1 \\ \prec A_1 \vee A_2 \\ \prec \neg A_1 \vee A_2 \\ \prec \neg A_1 \vee A_4 \vee A_3 \\ \prec \neg A_1 \vee \neg A_4 \vee A_3 \\ \prec \neg A_5 \vee A_5 \end{array}$$


Properties of the Clause Ordering

Proposition 3.16

1. The orderings on literals and clauses are total and well-founded.
2. Let C and D be clauses with $A = \max(C)$, $B = \max(D)$, where $\max(C)$ denotes the maximal atom in C .
 - (i) If $A \succ B$ then $C \succ D$.
 - (ii) If $A = B$, A occurs negatively in C but only positively in D , then $C \succ D$.

Stratified Structure of Clause Sets

Let $A \succ B$. Clause sets are then stratified in this form:



Closure of Clause Sets under Res

$$\begin{aligned}
 Res(N) &= \{C \mid C \text{ is concl. of a rule in } Res \text{ w/ premises in } N\} \\
 Res^0(N) &= N \\
 Res^{n+1}(N) &= Res(Res^n(N)) \cup Res^n(N), \text{ for } n \geq 0 \\
 Res^*(N) &= \bigcup_{n \geq 0} Res^n(N)
 \end{aligned}$$

N is called *saturated* (w. r. t. resolution), if $Res(N) \subseteq N$.

Proposition 3.17

- (i) $Res^*(N)$ is saturated.
- (ii) Res is refutationally complete, iff for each set N of ground clauses:

$$N \models \perp \Leftrightarrow \perp \in Res^*(N)$$

Construction of Interpretations

Given: set N of ground clauses, atom ordering \succ .

Wanted: Herbrand interpretation I such that

- “many” clauses from N are valid in I ;
- $I \models N$, if N is saturated and $\perp \notin N$.

Construction according to \succ , starting with the minimal clause.

Main Ideas of the Construction

- Clauses are considered in the order given by \prec .
- When considering C , one already has a partial interpretation I_C (initially $I_C = \emptyset$) available.
- If C is true in the partial interpretation I_C , nothing is done. ($\Delta_C = \emptyset$).
- If C is false, one would like to change I_C such that C becomes true.
- Changes should, however, be *monotone*. One never deletes anything from I_C and the truth value of clauses smaller than C should be maintained the way it was in I_C .
- Hence, one chooses $\Delta_C = \{A\}$ if, and only if, C is false in I_C , if A occurs positively in C (*adding A will make C become true*) and if this occurrence in C is strictly maximal in the ordering on literals (*changing the truth value of A has no effect on smaller clauses*).

Construction of Candidate Interpretations

Let N, \succ be given. We define sets I_C and Δ_C for all ground clauses C over the given signature inductively over \succ :

$$I_C := \bigcup_{C \succ D} \Delta_D$$
$$\Delta_C := \begin{cases} \{A\}, & \text{if } C \in N, C = C' \vee A, A \succ C', I_C \not\models C \\ \emptyset, & \text{otherwise} \end{cases}$$

We say that C produces A , if $\Delta_C = \{A\}$.

The *candidate interpretation* for N (w. r. t. \succ) is given as $I_N^\succ := \bigcup_C \Delta_C$. (We also simply write I_N or I for I_N^\succ if \succ is either irrelevant or known from the context.)

Example


Let $A_5 \succ A_4 \succ A_3 \succ A_2 \succ A_1 \succ A_0$ (max. literals in red)

	clauses C	I_C	Δ_C	Remarks
1	$\neg A_0$	\emptyset	\emptyset	true in I_C
2	$A_0 \vee A_1$	\emptyset	$\{A_1\}$	A_1 maximal
3	$A_1 \vee A_2$	$\{A_1\}$	\emptyset	true in I_C
4	$\neg A_1 \vee A_2$	$\{A_1\}$	$\{A_2\}$	A_2 maximal
5	$\neg A_1 \vee A_4 \vee A_3 \vee A_0$	$\{A_1, A_2\}$	$\{A_4\}$	A_4 maximal
6	$\neg A_1 \vee \neg A_4 \vee A_3$	$\{A_1, A_2, A_4\}$	\emptyset	A_3 not maximal; <i>min. counter-ex.</i>
7	$\neg A_1 \vee A_5$	$\{A_1, A_2, A_4\}$	$\{A_5\}$	

$I = \{A_1, A_2, A_4, A_5\}$ is not a model of the clause set
 \Rightarrow there exists a *counterexample*.

Structure of N, \succ

Let $A \succ B$; producing a new atom does not affect smaller clauses.



Some Properties of the Construction

Proposition 3.18

- (i) $C = \neg A \vee C' \Rightarrow$ no $D \succeq C$ produces A .
- (ii) C productive $\Rightarrow I_C \cup \Delta_C \models C$.
- (iii) Let $D' \succ D \succeq C$. Then

$$I_D \cup \Delta_D \models C \Rightarrow I_{D'} \cup \Delta_{D'} \models C \text{ and } I_N \models C.$$

If, in addition, $C \in N$ or $\max(D) \succ \max(C)$:

$$I_D \cup \Delta_D \not\models C \Rightarrow I_{D'} \cup \Delta_{D'} \not\models C \text{ and } I_N \not\models C.$$

(iv) Let $D' \succ D \succ C$. Then

$$I_D \models C \Rightarrow I_{D'} \models C \text{ and } I_N \models C.$$

If, in addition, $C \in N$ or $\max(D) \succ \max(C)$:

$$I_D \not\models C \Rightarrow I_{D'} \not\models C \text{ and } I_N \not\models C.$$

(v) $D = C \vee A$ produces $A \Rightarrow I_N \not\models C$.

Resolution Reduces Counterexamples

$$\frac{\neg A_1 \vee A_4 \vee A_3 \vee A_0 \quad \neg A_1 \vee \neg A_4 \vee A_3}{\neg A_1 \vee \neg A_1 \vee A_3 \vee A_3 \vee A_0}$$

Construction of I for the extended clause set:

clauses C	I_C	Δ_C	Remarks
$\neg A_0$	\emptyset	\emptyset	
$A_0 \vee A_1$	\emptyset	$\{A_1\}$	
$A_1 \vee A_2$	$\{A_1\}$	\emptyset	
$\neg A_1 \vee A_2$	$\{A_1\}$	$\{A_2\}$	
$\neg A_1 \vee \neg A_1 \vee A_3 \vee A_3 \vee A_0$	$\{A_1, A_2\}$	\emptyset	A_3 occurs twice <i>minimal counter-ex.</i>
$\neg A_1 \vee A_4 \vee A_3 \vee A_0$	$\{A_1, A_2\}$	$\{A_4\}$	
$\neg A_1 \vee \neg A_4 \vee A_3$	$\{A_1, A_2, A_4\}$	\emptyset	counterexample
$\neg A_1 \vee A_5$	$\{A_1, A_2, A_4\}$	$\{A_5\}$	

The same I , but smaller counterexample, hence some progress was made.

Factorization Reduces Counterexamples

$$\frac{\neg A_1 \vee \neg A_1 \vee A_3 \vee A_3 \vee A_0}{\neg A_1 \vee \neg A_1 \vee A_3 \vee A_0}$$

Construction of I for the extended clause set:

clauses C	I_C	Δ_C	Remarks
$\neg A_0$	\emptyset	\emptyset	
$A_0 \vee A_1$	\emptyset	$\{A_1\}$	
$A_1 \vee A_2$	$\{A_1\}$	\emptyset	
$\neg A_1 \vee A_2$	$\{A_1\}$	$\{A_2\}$	
$\neg A_1 \vee \neg A_1 \vee A_3 \vee A_0$	$\{A_1, A_2\}$	$\{A_3\}$	
$\neg A_1 \vee \neg A_1 \vee A_3 \vee A_3 \vee A_0$	$\{A_1, A_2, A_3\}$	\emptyset	true in I_C
$\neg A_1 \vee A_4 \vee A_3 \vee A_0$	$\{A_1, A_2, A_3\}$	\emptyset	
$\neg A_1 \vee \neg A_4 \vee A_3$	$\{A_1, A_2, A_3\}$	\emptyset	true in I_C
$\neg A_3 \vee A_5$	$\{A_1, A_2, A_3\}$	$\{A_5\}$	

The resulting $I = \{A_1, A_2, A_3, A_5\}$ is a model of the clause set.

Model Existence Theorem

Theorem 3.19 (Bachmair & Ganzinger 1990) *Let \succ be a clause ordering, let N be saturated w. r. t. Res, and suppose that $\perp \notin N$. Then $I_N^\succ \models N$.*

Corollary 3.20 *Let N be saturated w. r. t. Res. Then $N \models \perp \Leftrightarrow \perp \in N$.*

Proof of Theorem 3.19. Suppose $\perp \notin N$, but $I_N^\succ \not\models N$. Let $C \in N$ minimal (in \succ) such that $I_N^\succ \not\models C$. Since C is false in I_N , C is not productive. As $C \neq \perp$ there exists a maximal atom A in C .

Case 1: $C = \neg A \vee C'$ (i. e., the maximal atom occurs negatively)

$\Rightarrow I_N \models A$ and $I_N \not\models C'$

\Rightarrow some $D = D' \vee A \in N$ produces A . As $\frac{D' \vee A \quad \neg A \vee C'}{D' \vee C'}$, we infer that $D' \vee C' \in N$, and $C \succ D' \vee C'$ and $I_N \not\models D' \vee C'$

\Rightarrow contradicts minimality of C .

Case 2: $C = C' \vee A \vee A$. Then $\frac{C' \vee A \vee A}{C' \vee A}$ yields a smaller counterexample $C' \vee A \in N$. \Rightarrow contradicts minimality of C . \square

Compactness of Propositional Logic

Theorem 3.21 (Compactness) *Let N be a set of propositional formulas. Then N is unsatisfiable, if and only if some finite subset $M \subseteq N$ is unsatisfiable.*

Proof. “ \Leftarrow ”: trivial.

“ \Rightarrow ”: Let N be unsatisfiable.

$\Rightarrow Res^*(N)$ unsatisfiable

$\Rightarrow \perp \in Res^*(N)$ by refutational completeness of resolution