

6.3 Reduction Pairs and Argument Filterings

Goal: Show the non-existence of K -minimal infinite rewrite sequences

$$t_1 \rightarrow_R^* u_1 \rightarrow_K t_2 \rightarrow_R^* u_2 \rightarrow_K \dots$$

using well-founded orderings.

We observe that the requirements for the orderings used here are less restrictive than for reduction orderings:

K -rules are only used at the top, so we need stability under substitutions, but compatibility with contexts is unnecessary.

While \rightarrow_K -steps should be decreasing, for \rightarrow_R -steps it would be sufficient to show that they are not increasing.

This motivates the following definitions:

Rewrite quasi-ordering \succsim :

reflexive and transitive binary relation, stable under substitutions, compatible with contexts.

Reduction pair (\succsim, \succ) :

\succsim is a rewrite quasi-ordering.

\succ is a well-founded ordering that is stable under substitutions.

\succsim and \succ are compatible: $\succsim \circ \succ \subseteq \succ$ or $\succ \circ \succsim \subseteq \succ$.

(In practice, \succ is almost always the strict part of the quasi-ordering \succsim .)

Clearly, for any reduction ordering \succ , (\succ, \succ) is a reduction pair. More general reduction pairs can be obtained using argument filterings:

Argument filtering π :

$$\pi : \Omega \cup \Omega^\# \rightarrow \mathbb{N} \cup \text{list of } \mathbb{N}$$

$$\pi(f) = \begin{cases} i \in \{1, \dots, \text{arity}(f)\}, \text{ or} \\ [i_1, \dots, i_k], \text{ where } 1 \leq i_1 < \dots < i_k \leq \text{arity}(f), 0 \leq k \leq \text{arity}(f) \end{cases}$$

Extension to terms:

$$\pi(x) = x$$

$$\pi(f(t_1, \dots, t_n)) = \pi(t_i), \text{ if } \pi(f) = i$$

$$\pi(f(t_1, \dots, t_n)) = f'(\pi(t_{i_1}), \dots, \pi(t_{i_k})), \text{ if } \pi(f) = [i_1, \dots, i_k],$$

where f'/k is a new function symbol.

Let \succ be a reduction ordering, let π be an argument filtering. Define $s \succ_{\pi} t$ iff $\pi(s) \succ \pi(t)$ and $s \succeq_{\pi} t$ iff $\pi(s) \succeq \pi(t)$.

Lemma 6.2 $(\succeq_{\pi}, \succ_{\pi})$ is a reduction pair.

Proof. Follows from the following two properties:

$\pi(s\sigma) = \pi(s)\sigma_{\pi}$, where σ_{π} is the substitution that maps x to $\pi(\sigma(x))$.

$$\pi(s[u]_p) = \begin{cases} \pi(s), & \text{if } p \text{ does not correspond to any position in } \pi(s) \\ \pi(s)[\pi(u)]_q, & \text{if } p \text{ corresponds to } q \text{ in } \pi(s) \end{cases} \quad \square$$

For interpretation-based orderings (such as polynomial orderings) the idea of “cutting out” certain subterms can be included directly in the definition of the ordering:

Reduction pairs by interpretation:

Let \mathcal{A} be a Σ -algebra; let \succ be a well-founded strict partial ordering on its universe.

Assume that all interpretations $f_{\mathcal{A}}$ of function symbols are *weakly monotone*, i. e., $a_i \succeq b_i$ implies $f(a_1, \dots, a_n) \succeq f(b_1, \dots, b_n)$ for all $a_i, b_i \in U_{\mathcal{A}}$.

Define $s \succeq_{\mathcal{A}} t$ iff $\mathcal{A}(\beta)(s) \succeq \mathcal{A}(\beta)(t)$ for all assignments $\beta : X \rightarrow U_{\mathcal{A}}$; define $s \succ_{\mathcal{A}} t$ iff $\mathcal{A}(\beta)(s) \succ \mathcal{A}(\beta)(t)$ for all assignments $\beta : X \rightarrow U_{\mathcal{A}}$.

Then $(\succeq_{\mathcal{A}}, \succ_{\mathcal{A}})$ is a reduction pair.

For polynomial orderings, this definition permits interpretations of function symbols where some variable does not occur at all (e. g., $P_f(X, Y) = 2X + 1$ for a *binary* function symbol). It is no longer required that every variable must occur with some positive coefficient.

Theorem 6.3 (Arts and Giesl) *Let K be a cycle in the dependency graph of the TRS R . If there is a reduction pair (\succeq, \succ) such that*

- $l \succeq r$ for all $l \rightarrow r \in R$,
- $l \succeq r$ or $l \succ r$ for all $l \rightarrow r \in K$,
- $l \succ r$ for at least one $l \rightarrow r \in K$,

then there is no K -minimal infinite sequence.

Proof. Assume that

$$t_1 \rightarrow_R^* u_1 \rightarrow_K t_2 \rightarrow_R^* u_2 \rightarrow_K \dots$$

is a K -minimal infinite rewrite sequence.

As $l \succsim r$ for all $l \rightarrow r \in R$, we obtain $t_i \succsim u_i$ by stability under substitutions, compatibility with contexts, reflexivity and transitivity.

As $l \succsim r$ or $l \succ r$ for all $l \rightarrow r \in K$, we obtain $u_i (\succsim \cup \succ) t_{i+1}$ by stability under substitutions.

So we get an infinite $(\succsim \cup \succ)$ -sequence containing infinitely many \succ -steps (since every DP in K , in particular the one for which $l \succ r$ holds, is used infinitely often).

By compatibility of \succsim and \succ , we can transform this into an infinite \succ -sequence, contradicting well-foundedness. \square

The idea can be extended to SCCs in the same way as for the subterm criterion:

Search for a reduction pair (\succsim, \succ) such that $l \succsim r$ for all $l \rightarrow r \in R$ and $l \succsim r$ or $l \succ r$ for all DPs $l \rightarrow r$ in the SCC. Delete all DPs in the SCC for which $l \succ r$. Then re-compute SCCs for the remaining graph and re-start.

Example: Consider the following TRS R from [Arts and Giesl]:

$$\text{minus}(x, 0) \rightarrow x \tag{1}$$

$$\text{minus}(s(x), s(y)) \rightarrow \text{minus}(x, y) \tag{2}$$

$$\text{quot}(0, s(y)) \rightarrow 0 \tag{3}$$

$$\text{quot}(s(x), s(y)) \rightarrow s(\text{quot}(\text{minus}(x, y), s(y))) \tag{4}$$

(R is not contained in any simplification ordering, since the left-hand side of rule (4) is embedded in the right-hand side after instantiating y by $s(x)$.)


R has three dependency pairs:

$$\text{minus}^\sharp(s(x), s(y)) \rightarrow \text{minus}^\sharp(x, y) \tag{5}$$

$$\text{quot}^\sharp(s(x), s(y)) \rightarrow \text{quot}^\sharp(\text{minus}(x, y), s(y)) \tag{6}$$

$$\text{quot}^\sharp(s(x), s(y)) \rightarrow \text{minus}^\sharp(x, y) \tag{7}$$

The dependency graph of R is



There are exactly two SCCs (and also two cycles). The cycle at (5) can be handled using the subterm criterion with $\pi(\mathit{minus}^\sharp) = 1$. For the cycle at (6) we can use an argument filtering π that maps minus to 1 and leaves all other function symbols unchanged (that is, $\pi(g) = [1, \dots, \mathit{arity}(g)]$ for every g different from minus .) After applying the argument filtering, we compare left and right-hand sides using an LPO with precedence $\mathit{quot} > s$ (the precedence of other symbols is irrelevant). We obtain $l \succ r$ for (6) and $l \succsim r$ for (1), (2), (3), (4), so the previous theorem can be applied.

The methods described so far are particular cases of *DP processors*:

A DP processor

$$\frac{(G, R)}{(G_1, R_1), \dots, (G_n, R_n)}$$

takes a graph G and a TRS R as input and produces a set of pairs consisting of a graph and a TRS.

It is sound and complete if there are K -minimal infinite sequences for G and R if and only if there are K -minimal infinite sequences for at least one of the pairs (G_i, R_i) .

Examples:

$$\frac{(G, R)}{(SCC_1, R), \dots, (SCC_n, R)}$$

where SCC_1, \dots, SCC_n are the strongly connected components of G .

$$\frac{(G, R)}{(G \setminus N, R)}$$

if there is an SCC of G and a simple projection π such that $\pi(l) \succeq \pi(r)$ for all DPs $l \rightarrow r$ in the SCC, and N is the set of DPs of the SCC for which $\pi(l) \triangleright \pi(r)$.

(and analogously for reduction pairs)

The dependency method can also be used for proving termination of *innermost rewriting*: $s \xrightarrow{i}_R t$ if $s \rightarrow_R t$ at position p and no rule of R can be applied at a position strictly below p . (DP processors for innermost termination are more powerful than for ordinary termination, and for program analysis, innermost termination is usually sufficient.)