

Topic IV: Tensors

Discrete Topics in Data Mining
Universität des Saarlandes, Saarbrücken
Winter Semester 2012/13

Topic IV: Tensors

- 1. What is a ... tensor?**
- 2. Basic Operations**
- 3. Tensor Decompositions and Rank**
 - 3.1. CP Decomposition**
 - 3.2. Tensor Rank**
 - 3.3. Tucker Decomposition**

I admire the elegance of your method of computation; it must be nice to ride through these fields upon the horse of true mathematics while the like of us have to make our way laboriously on foot.

Albert Einstein
in a letter to Tullio Levi-Civita

What is a ... tensor?

- A **tensor** is a multi-way extension of a matrix
 - A multi-dimensional array
 - A multi-linear map
- In particular, the following are all tensors:

What is a ... tensor?

- A **tensor** is a multi-way extension of a matrix
 - A multi-dimensional array
 - A multi-linear map
- In particular, the following are all tensors:
 - Scalars

13

What is a ... tensor?

- A **tensor** is a multi-way extension of a matrix
 - A multi-dimensional array
 - A multi-linear map
- In particular, the following are all tensors:
 - Scalars
 - Vectors

$(13, 42, 2011)$

What is a ... tensor?

- A **tensor** is a multi-way extension of a matrix
 - A multi-dimensional array
 - A multi-linear map
- In particular, the following are all tensors:
 - Scalars
 - Vectors
 - Matrices

$$\begin{pmatrix} 8 & 1 & 6 \\ 3 & 5 & 7 \\ 4 & 9 & 2 \end{pmatrix}$$


What is a ... tensor?

- A **tensor** is a multi-way extension of a matrix
 - A multi-dimensional array
 - A multi-linear map
- In particular, the following are all tensors:
 - Scalars
 - Vectors
 - Matrices



What is a ... tensor?


- A **tensor** is a multi-way extension of a matrix
 - A multi-dimensional array
 - A multi-linear map
- In particular, the following are all tensors:
 - Scalars
 - Vectors
 - Matrices




Why Tensors?

- Tensors can be used when matrices are not enough
- A matrix can represent a binary relation
 - A tensor can represent an n -ary relation
 - E.g. subject–predicate–object data
 - A tensor can represent a set of binary relations
 - Or other matrices
- A matrix can represent a matrix
 - A tensor can represent a series/set of matrices
 - But using tensors for time series should be approached with care


Fibres and Slices




(a) Mode-1 (column) fibers: $\mathbf{x}_{:jk}$




(b) Mode-2 (row) fibers: $\mathbf{x}_{i:k}$




(c) Mode-3 (tube) fibers: $\mathbf{x}_{ij:}$



(a) Horizontal slices: $\mathbf{X}_{i::}$



(b) Lateral slices: $\mathbf{X}_{:j:}$



(c) Frontal slices: $\mathbf{X}_{::k}$ (or \mathbf{X}_k)

Kolda & Bader 2009

Basic Operations

- Tensors require extensions to the standard linear algebra operations for matrices
- A **multi-way vector outer product** is a tensor where each element is the product of corresponding elements in vectors: $\mathcal{X} = \mathbf{a} \circ \mathbf{b} \circ \mathbf{c}$, $(\mathcal{X})_{ijk} = a_i b_j c_k$
- A **tensor inner product** of two same-sized tensors is the sum of the element-wise products of their values:
$$\langle \mathcal{X}, \mathcal{Y} \rangle = \sum_{i=1}^I \sum_{j=1}^J \cdots \sum_{z=1}^Z x_{ij \dots z} y_{ij \dots z}$$

Tensor Matricization

- Tensor **matricization** unfolds an N -way tensor into a matrix
 - **Mode- n matricization** arranges the mode- n fibers as columns of a matrix
 - Denoted $\mathbf{X}_{(n)}$
 - As many rows as is the dimensionality of the n th mode
 - As many columns as is the product of the dimensions of the other modes
- If \mathcal{X} is an N -way tensor of size $I_1 \times I_2 \times \dots \times I_N$, then $\mathbf{X}_{(n)}$ maps element x_{i_1, i_2, \dots, i_N} into (i_n, j) where
$$j = 1 + \sum_{k=1}^N (i_k - 1) J_k [k \neq n] \text{ with } J_k = \prod_{m=1}^{k-1} I_m [m \neq n]$$

Matricization Example

$$\mathbf{x} = \begin{pmatrix} 0 & 1 \\ 1 & 0 \\ 0 & 1 \end{pmatrix}$$

Matricization Example

$$\mathcal{X} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

Matricization Example

$$\mathbf{x} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

$$\mathbf{X}_{(1)} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 \end{pmatrix}$$

Matricization Example

$$\mathbf{x} = \begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix} \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix}$$

$$\mathbf{X}_{(1)} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 \end{pmatrix}$$

$$\mathbf{X}_{(2)} = \begin{pmatrix} 1 & 0 & 0 & -1 \\ 0 & 1 & 1 & 0 \end{pmatrix}$$

$$\mathbf{X}_{(3)} = \begin{pmatrix} 1 & 0 & 0 & 1 \\ 0 & 1 & -1 & 0 \end{pmatrix}$$

Tensor Multiplication

- Let \mathcal{X} be an N -way tensor of size $I_1 \times I_2 \times \dots \times I_N$, and let \mathbf{U} be a matrix of size $J \times I_n$
 - The **n -mode matrix product** of \mathcal{X} with \mathbf{U} , $\mathcal{X} \times_n \mathbf{U}$ is of size $I_1 \times I_2 \times \dots \times I_{n-1} \times J \times I_{n+1} \times \dots \times I_N$
 - $(\mathcal{X} \times_n \mathbf{U})_{i_1 \dots i_{n-1} j i_{n+1} \dots i_N} = \sum_{i_n=1}^{I_n} x_{i_1 i_2 \dots i_N} u_{j i_n}$
 - Each mode- n fibre is multiplied by the matrix \mathbf{U}
 - In terms of unfold tensors: $\mathcal{Y} = \mathcal{X} \times_n \mathbf{U} \iff \mathbf{Y}_{(n)} = \mathbf{U} \mathbf{X}_{(n)}$
- The **n -mode vector product** is denoted $\mathcal{X} \bar{\times}_n \mathbf{v}$
 - The result is of order $N-1$
 - $(\mathcal{X} \bar{\times}_n \mathbf{v})_{i_1 \dots i_{n-1} i_{n+1} \dots i_N} = \sum_{i_n=1}^{I_n} x_{i_1 i_2 \dots i_N} v_{i_n}$
 - Inner product between mode- n fibres and vector \mathbf{v}

Kronecker Matrix Product

- Element-per-matrix product
- n -by- m and j -by- k matrices give nj -by- mk matrix

$$\mathbf{A} \otimes \mathbf{B} = \begin{pmatrix} a_{1,1}\mathbf{B} & a_{1,2}\mathbf{B} & \cdots & a_{1,m}\mathbf{B} \\ a_{2,1}\mathbf{B} & a_{2,2}\mathbf{B} & \cdots & a_{2,m}\mathbf{B} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n,1}\mathbf{B} & a_{n,2}\mathbf{B} & \cdots & a_{n,m}\mathbf{B} \end{pmatrix}$$

Khatri–Rao Matrix Product

- Element-per-column product
 - Number of columns must match
- n -by- m and k -by- m matrices give nk -by- m matrix

$$\mathbf{A} \odot \mathbf{B} = \begin{pmatrix} a_{1,1}\mathbf{b}_1 & a_{1,2}\mathbf{b}_2 & \cdots & a_{1,m}\mathbf{b}_m \\ a_{2,1}\mathbf{b}_1 & a_{2,2}\mathbf{b}_2 & \cdots & a_{2,m}\mathbf{b}_m \\ \vdots & \vdots & \ddots & \vdots \\ a_{n,1}\mathbf{b}_1 & a_{n,2}\mathbf{b}_2 & \cdots & a_{n,m}\mathbf{b}_m \end{pmatrix}$$

Hadamard Matrix Product

- The element-wise matrix product
- Two matrices of size n -by- m , resulting matrix of size n -by- m

$$\mathbf{A} * \mathbf{B} = \begin{pmatrix} a_{1,1}b_{1,1} & a_{1,2}b_{1,2} & \cdots & a_{1,m}b_{1,m} \\ a_{2,1}b_{2,1} & a_{2,2}b_{2,2} & \cdots & a_{2,m}b_{2,m} \\ \vdots & \vdots & \ddots & \vdots \\ a_{n,1}b_{n,1} & a_{n,2}b_{n,2} & \cdots & a_{n,m}b_{n,m} \end{pmatrix}$$

Some identities

$$(\mathbf{A} \otimes \mathbf{B})(\mathbf{C} \otimes \mathbf{D}) = \mathbf{AC} \otimes \mathbf{BD}$$

$$(\mathbf{A} \otimes \mathbf{B})^\dagger = \mathbf{A}^\dagger \otimes \mathbf{B}^\dagger$$

$$\mathbf{A} \odot \mathbf{B} \odot \mathbf{C} = (\mathbf{A} \odot \mathbf{B}) \odot \mathbf{C} = \mathbf{A} \odot (\mathbf{B} \odot \mathbf{C})$$

$$(\mathbf{A} \odot \mathbf{B})^T (\mathbf{A} \odot \mathbf{B}) = \mathbf{A}^T \mathbf{A} * \mathbf{B}^T \mathbf{B}$$

$$(\mathbf{A} \odot \mathbf{B})^\dagger = \left((\mathbf{A}^T \mathbf{A}) * (\mathbf{B}^T \mathbf{B}) \right)^\dagger (\mathbf{A} \odot \mathbf{B})^T$$

\mathbf{A}^\dagger is the Moore–Penrose pseudo-inverse

Tensor Decompositions and Rank


- A matrix decomposition represents the given matrix as a product of two (or more) **factor matrices**
- The **rank** of a matrix \mathbf{M} is the
 - Number of linearly independent rows (*row rank*)
 - Number of linearly independent columns (*column rank*)
 - Number of rank-1 matrices needed to be summed to get \mathbf{M} (*Schein rank*)
 - Rank-1 matrix is an outer product of two vectors
 - They all are equivalent

Tensor Decompositions and Rank

- A matrix decomposition represents the given matrix as a product of two (or more) **factor matrices**
- The **rank** of a matrix \mathbf{M} is the
 - Number of linearly independent rows (*row rank*)
 - Number of linearly independent columns (*column rank*)
 - Number of rank-1 matrices needed to be summed to get \mathbf{M} (*Schein rank*)
 - Rank-1 matrix is an outer product of two vectors
 - They all are equivalent


This we generalize

Rank-1 Tensors




$$\mathbf{X} = \mathbf{a} \circ \mathbf{b}$$

Rank-1 Tensors



$$\mathcal{X} = \mathbf{a} \circ \mathbf{b} \circ \mathbf{c}$$

The CP Tensor Decomposition



$$x_{ijk} \approx \sum_{r=1}^R a_{ir} b_{jr} c_{kr}$$

More on CP

- The *size* of the CP factorization is the number of rank-1 tensors involved
- The factorization can also be written using N factor matrix (for order- N tensor)
 - All column vectors are collected in one matrix, all row vectors in other, all tube vectors in third, etc.
 - These matrices are typically called **A**, **B**, and **C** for 3rd order tensors

CANDECOM, PARAFAC, ...

Name	Proposed by
Polyadic Form of a Tensor	Hitchcock, 1927 [105]
PARAFAC (Parallel Factors)	Harshman, 1970 [90]
CANDECOMP or CAND (Canonical decomposition)	Carroll and Chang, 1970 [38]
Topographic Components Model	Möcks, 1988 [166]
CP (CANDECOMP/PARAFAC)	Kiers, 2000 [122]

Table 3.1: Some of the many names for the CP decomposition.

Another View on the CP

- Using matricization, we can re-write the CP decomposition
 - One equation per mode

$$\mathbf{X}_{(1)} = \mathbf{A}(\mathbf{C} \odot \mathbf{B})^T$$

$$\mathbf{X}_{(2)} = \mathbf{B}(\mathbf{C} \odot \mathbf{A})^T$$

$$\mathbf{X}_{(3)} = \mathbf{C}(\mathbf{B} \odot \mathbf{A})^T$$

Solving CP: The ALS Approach

1. Fix **B** and **C** and solve **A**
2. Solve **B** and **C** similarly
3. Repeat until convergence

Solving CP: The ALS Approach

1. Fix **B** and **C** and solve **A**
2. Solve **B** and **C** similarly
3. Repeat until convergence

$$\min_{\mathbf{A}} \left\| \mathbf{X}_{(1)} - \mathbf{A}(\mathbf{C} \odot \mathbf{B})^T \right\|_F$$

Solving CP: The ALS Approach

1. Fix **B** and **C** and solve **A**
2. Solve **B** and **C** similarly
3. Repeat until convergence

$$\min_{\mathbf{A}} \left\| \mathbf{X}_{(1)} - \mathbf{A}(\mathbf{C} \odot \mathbf{B})^T \right\|_F$$

$$\mathbf{A} = \mathbf{X}_{(1)} \left((\mathbf{C} \odot \mathbf{B})^T \right)^\dagger$$

Solving CP: The ALS Approach

1. Fix **B** and **C** and solve **A**
2. Solve **B** and **C** similarly
3. Repeat until convergence

$$\min_{\mathbf{A}} \left\| \mathbf{X}_{(1)} - \mathbf{A}(\mathbf{C} \odot \mathbf{B})^T \right\|_F$$

$$\mathbf{A} = \mathbf{X}_{(1)} \left((\mathbf{C} \odot \mathbf{B})^T \right)^\dagger$$

$$\mathbf{A} = \mathbf{X}_{(1)} (\mathbf{C} \odot \mathbf{B}) (\mathbf{C}^T \mathbf{C} * \mathbf{B}^T \mathbf{B})^\dagger$$

Solving CP: The ALS Approach

1. Fix **B** and **C** and solve **A**
2. Solve **B** and **C** similarly
3. Repeat until convergence

$$\min_{\mathbf{A}} \left\| \mathbf{X}_{(1)} - \mathbf{A}(\mathbf{C} \odot \mathbf{B})^T \right\|_F$$


$$\mathbf{A} = \mathbf{X}_{(1)} \left((\mathbf{C} \odot \mathbf{B})^T \right)^\dagger$$

$$\mathbf{A} = \mathbf{X}_{(1)} (\mathbf{C} \odot \mathbf{B}) (\mathbf{C}^T \mathbf{C} * \mathbf{B}^T \mathbf{B})^\dagger$$

R-by-R matrix

Tensor Rank

- The **rank** of a tensor is the minimum number of rank-1 tensors needed to represent the tensor exactly
 - The CP decomposition of size R
 - Generalizes the matrix Schein rank



Tensor Rank Oddities #1

- The rank of a (real-valued) tensor is different over reals and over complex numbers.
 - With reals, the rank can be *larger* than the largest dimension
 - $\text{rank}(\mathcal{X}) \leq \min\{IJ, IK, JK\}$ for I -by- J -by- K tensor

$$\mathcal{X} = \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right)$$

$$\mathbf{A} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ -i & i \end{pmatrix},$$

$$\mathbf{B} = \frac{1}{\sqrt{2}} \begin{pmatrix} 1 & 1 \\ i & -i \end{pmatrix},$$

$$\mathbf{C} = \begin{pmatrix} 1 & 1 \\ i & -1 \end{pmatrix}$$

Tensor Rank Oddities #1

- The rank of a (real-valued) tensor is different over reals and over complex numbers.
 - With reals, the rank can be *larger* than the largest dimension
 - $\text{rank}(\mathcal{X}) \leq \min\{IJ, IK, JK\}$ for I -by- J -by- K tensor

$$\mathcal{X} = \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right)$$
$$\mathbf{A} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \end{pmatrix},$$
$$\mathbf{B} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix},$$
$$\mathbf{C} = \begin{pmatrix} 1 & 1 & 0 \\ -1 & 1 & 1 \end{pmatrix}$$

Tensor Rank Oddities #1

- The rank of a (real-valued) tensor is different over reals and over complex numbers.
 - With reals, the rank can be *larger* than the largest dimension

$$\mathcal{X} = \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right)$$
$$\mathbf{A} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \end{pmatrix},$$
$$\mathbf{B} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix},$$
$$\mathbf{C} = \begin{pmatrix} 1 & 1 & 0 \\ -1 & 1 & 1 \end{pmatrix}$$

Tensor Rank Oddities #1

- The rank of a (real-valued) tensor is different over reals and over complex numbers.
 - With reals, the rank can be *larger* than the largest dimension
 - $\text{rank}(\mathcal{X}) \leq \min\{IJ, IK, JK\}$ for I -by- J -by- K tensor

$$\mathcal{X} = \left(\begin{pmatrix} 1 & 0 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 0 & 1 \\ -1 & 0 \end{pmatrix} \right)$$
$$\mathbf{A} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & -1 \end{pmatrix},$$
$$\mathbf{B} = \begin{pmatrix} 1 & 0 & 1 \\ 0 & 1 & 1 \end{pmatrix},$$
$$\mathbf{C} = \begin{pmatrix} 1 & 1 & 0 \\ -1 & 1 & 1 \end{pmatrix}$$


Tensor Rank Oddities #2

- There are tensors of rank R that can be approximated arbitrarily well with tensors of rank R' for some $R' < R$.
 - That is, there are no *best* low-rank approximation for such tensors.
 - Eckart–Young-theorem shows this is impossible with matrices.
 - The smallest such R' is called the **border rank** of the tensor.

Tensor Rank Oddities #3

- The rank- R CP decomposition of a rank- R tensor is *essentially unique* under mild conditions.
 - Essentially unique = only scaling and permuting are allowed.
 - Does not contradict #2, as this is the rank decomposition, not low-rank decomposition.
 - Again, not true for matrices (unless orthogonality etc. is required).

The Tucker Tensor Decomposition



$$x_{ijk} \approx \sum_{p=1}^P \sum_{q=1}^Q \sum_{r=1}^R g_{pqr} a_{ip} b_{jq} c_{kr}$$

Tucker Decomposition

- Many degrees of freedom: often **A**, **B**, and **C** are required to be orthogonal
- If $P=Q=R$ and core tensor \mathcal{G} is hyper-diagonal, then Tucker decomposition reduces to CP decomposition
- ALS-style methods are typically used
 - The matricized forms are

$$\mathbf{X}_{(1)} = \mathbf{A}\mathbf{G}_{(1)}(\mathbf{C} \otimes \mathbf{B})^T$$

$$\mathbf{X}_{(2)} = \mathbf{B}\mathbf{G}_{(2)}(\mathbf{C} \otimes \mathbf{A})^T$$

$$\mathbf{X}_{(3)} = \mathbf{C}\mathbf{G}_{(3)}(\mathbf{B} \otimes \mathbf{A})^T$$

Higher-Order SVD (HOSVD)

- One method to compute the Tucker decomposition
 - Set \mathbf{A} as the leading P left singular vectors of $\mathbf{X}_{(1)}$
 - Set \mathbf{B} as the leading Q left singular vectors of $\mathbf{X}_{(2)}$
 - Set \mathbf{C} as the leading R left singular vectors of $\mathbf{X}_{(3)}$
 - Set tensor \mathcal{G} as $\mathcal{X} \times_1 \mathbf{A}^T \times_2 \mathbf{B}^T \times_3 \mathbf{C}^T$

Comments

- Tensors generalize matrices
- Many matrix concepts generalize as well
 - But some don't
 - And some behave very differently
- Compared to matrix decomposition methods, tensor algorithms are in their youth
 - Notwithstanding that Tucker did his work in 60's